

CARLESON MEASURES AND LOGVINENKO-SEREDA SETS ON COMPACT MANIFOLDS

JOAQUIM ORTEGA-CERDÀ AND BHARTI PRIDHNANI

ABSTRACT. Given a compact Riemannian manifold M of dimension $m \geq 2$, we study the space of functions of $L^2(M)$ generated by eigenfunctions of eigenvalues less than $L \geq 1$ associated to the Laplace-Beltrami operator on M . On these spaces we give a characterization of the Carleson measures and the Logvinenko-Sereda sets.

INTRODUCTION AND STATEMENT OF THE RESULTS

Let (M, g) be a smooth, connected, compact Riemannian manifold without boundary of dimension $m \geq 2$. Let dV be the volume element of M associated to the metric g_{ij} . Let Δ_M be the Laplacian on M associated to the metric g_{ij} . It is given in local coordinates by

$$\Delta_M f = \frac{1}{\sqrt{|g|}} \sum_{i,j} \frac{\partial}{\partial x_i} \left(\sqrt{|g|} g^{ij} \frac{\partial f}{\partial x_j} \right),$$

where $|g| = \det(g_{ij})$ and $(g^{ij})_{ij}$ is the inverse matrix of $(g_{ij})_{ij}$. Since M is compact, g_{ij} and all its derivatives are bounded and we assume that the metric g is non-singular at each point of M .

Since M is compact, the spectrum of the Laplacian is discrete and there is a sequence of eigenvalues

$$0 \leq \lambda_1 \leq \lambda_2 \leq \dots \rightarrow \infty$$

and an orthonormal basis ϕ_i of smooth real eigenfunctions of the Laplacian i.e. $\Delta_M \phi_i = -\lambda_i \phi_i$. So $L^2(M)$ decomposes into an orthogonal direct sum of eigenfunctions of the Laplacian.

We consider the following spaces of $L^2(M)$.

$$E_L = \left\{ f \in L^2(M) : f = \sum_{i=1}^{k_L} \beta_i \phi_i, \Delta_M \phi_i = -\lambda_i \phi_i, \lambda_{k_L} \leq L \right\},$$

Date: April 15, 2010.

2000 *Mathematics Subject Classification.* 35P99, 58C35, 58C40.

Supported by the project MTM2008-05561-C02-01 and the CIRIT grant 2005SGR00611.

where $L \geq 1$ and $k_L = \dim E_L$. E_L is the space of $L^2(M)$ generated by eigenfunctions of eigenvalues $\lambda \leq L$. Thus in E_L we consider functions in $L^2(M)$ with a restriction on the support of its Fourier transform. It is, in a sense, a Paley-Wiener type space on M with bandwidth L .

The motivation of this paper is to show that the spaces E_L behave like the space defined in $\mathbb{S}^d$ ($d > 1$) of spherical harmonics of degree less than $\sqrt{L}$. In fact, the space E_L is a generalization of the spherical harmonics and the role of them are played by the eigenfunctions. The cases $M = S^1$ and $M = \mathbb{S}^d$ ($d > 1$) have been studied in [14] and [11], respectively.

This similarity between eigenfunctions of the Laplacian and polynomials is not new, for instance, Donnelly and Fefferman in [1, Theorem 1], showed that on a compact manifold, an eigenfunction of eigenvalue λ behaves essentially like a polynomial of degree $\sqrt{\lambda}$. In this direction they proved the following local L^∞ Bernstein inequality.

Theorem (Donnelly-Fefferman). *Let M be as above. If u is an eigenfunction of the Laplacian $\Delta_M u = -\lambda u$, then there exists $r_0 = r_0(M)$ such that for all $r < r_0$ we have*

$$\max_{B(x,r)} |\nabla u| \leq \frac{C\lambda^{(m+2)/4}}{r} \max_{B(x,r)} |u|.$$

The proof of the above estimate is rather delicate. Donnelly and Fefferman conjectured that it is possible to replace $\lambda^{(m+2)/4}$ by $\sqrt{\lambda}$ in the inequality. If the conjecture holds, we have in particular, a global Bernstein type inequality:

$$(1) \quad \|\nabla u\|_\infty \lesssim \sqrt{\lambda} \|u\|_\infty.$$

In fact, this weaker estimate holds and a proof will be given later. This fact suggests that the right metric to study the space E_L should be rescaled by a factor $1/\sqrt{L}$ because in balls of radius $1/\sqrt{\lambda}$, a bounded eigenfunction of eigenvalue λ oscillates very little.

In the present work we will study for which measures $\mu = \{\mu_L\}_L$ one has

$$(2) \quad \int_M |f|^2 d\mu_L \approx \int_M |f|^2 dV, \quad \forall f \in E_L$$

with constants independent of f and L .

We will also study the inequality

$$\int_M |f|^2 d\mu_L \lesssim \int_M |f|^2 dV$$

that defines the Carleson measures and we will present a geometric characterization of them. Inequality (2) will be studied only for the special case $d\mu_L = \chi_{A_L} dV$, where $\mathcal{A} = \{A_L\}_L$ is a sequence of sets in

the manifold. In such case, when (2) holds, we say that $\mathcal{A}$ is a sequence of Logvinenko-Sereda sets. Our two main results are the following:

Theorem 1. *Let $\mu = \{\mu_L\}_L$ be a sequence of measures on M . Then μ is L^2 -Carleson for M if and only if there exists a $C > 0$ such that for all L*

$$\sup_{\xi \in M} \frac{\mu_L(B(\xi, 1/\sqrt{L}))}{\text{vol}(B(\xi, 1/\sqrt{L}))} \leq C.$$

Theorem 2. *The sequence of sets $\mathcal{A} = \{A_L\}_L$ is Logvinenko-Sereda if and only if there is an $r > 0$ such that*

$$\inf_L \inf_{z \in M} \frac{\text{vol}(A_L \cap B(z, r/\sqrt{L}))}{\text{vol}(B(z, r/\sqrt{L}))} > 0.$$

In what follows, when we write that $A \lesssim B$, $A \gtrsim B$ or $A \approx B$ we mean that there are constants depending only on the manifold such that $A \leq CB$, $A \geq CB$ or $C_1B \leq A \leq C_2B$, respectively. Also, the value of the constants appearing during a proof may change but they will be still denoted with the same letter. We will denote by $B(\xi, r)$ a geodesic ball in M of center ξ and radius r and $\mathbb{B}(z, r)$ will denote an Euclidean ball in $\mathbb{R}^m$ of center z and radius r .

The structure of the paper is the following: in the first section, we will explain the asymptotics of the reproducing kernel of the space E_L . In the second section, we shall discuss one of the tools used: the harmonic extension of functions in the space E_L . Following this, we will study the Carleson measures associated to M and we will prove Theorem 1. In the last section, Theorem 2 will be proved.

Acknowledgment. We thank professor M. Sanchón for valuable comments on the subject and professor S. Zelditch for providing us appropriate references.

1. THE REPRODUCING KERNEL OF E_L

Let

$$K_L(z, w) := \sum_{i=1}^{k_L} \phi_i(z) \phi_i(w) = \sum_{\lambda_i \leq L} \phi_i(z) \phi_i(w).$$

This function is the reproducing kernel of the space E_L , i.e.

$$\forall f \in E_L \quad f(z) = \langle f, K_L(z, \cdot) \rangle.$$

Note that $\dim(E_L) = k_L = \#\{\lambda_i \leq L\}$. The function K_L is also called the spectral function associated to the Laplacian. Hörmander in [5], proved the following estimates.

$$(1) \quad K_L(z, w) = O(L^{(m-1)/2}), \quad z \neq w.$$

$$(2) \quad K_L(z, z) = \frac{\sigma_m}{(2\pi)^m} L^{m/2} + O(L^{(m-1)/2}) \quad (\text{uniformly in } z \in M),$$

where $\sigma_m = 2\pi^{m/2}/(m\Gamma(m/2))$.

$$(3) \quad k_L = \frac{\text{vol}(M)\sigma_m}{(2\pi)^m} L^{m/2} + O(L^{(m-1)/2}).$$

In fact, in [5], there are estimates for the spectral function associated to any elliptic operator of order $n \geq 1$ with constants depending only on the manifold.

So, for L big enough we have $k_L \approx L^{m/2}$ and

$$\|K_L(z, \cdot)\|_2^2 = K_L(z, z) \approx L^{m/2} \approx k_L$$

with constants independent of L and z .

2. HARMONIC EXTENSION

In what follows, given $f \in E_L$, we will note by h the harmonic extension of f in $N = M \times \mathbb{R}$. The metric that we consider in N is the product metric, i.e. if we denote it by $\tilde{g}_{ij}$ for $i = 1, \dots, m+1$ then

$$(\tilde{g}_{ij})_{i,j=1,\dots,m+1} = \begin{pmatrix} (g_{ij})_{i,j=1}^m & 0 \\ 0 & 1 \end{pmatrix}.$$

Using this matrix, we can calculate the gradient and the Laplacian for N . If $h(z, t)$ is a function defined on N then

$$|\nabla_N h(z, t)|^2 = |\nabla_M h(z, t)|^2 + \left(\frac{\partial h}{\partial t}(z, t) \right)^2$$

and

$$\Delta_N h(z, t) = \Delta_M h(z, t) + \frac{\partial^2 h}{\partial t^2}(z, t).$$

Note that $|\nabla_M h(z, t)| \leq |\nabla_N h(z, t)|$.

Let $f \in E_L$, we know that

$$f = \sum_{i=1}^{k_L} \beta_i \phi_i, \quad \Delta_M \phi_i = -\lambda_i \phi_i, \quad 0 \leq \lambda_i \leq L.$$

Define for $(z, t) \in N$

$$h(z, t) = \sum_{i=1}^{k_L} \beta_i \phi_i(z) e^{\sqrt{\lambda_i} t}.$$

Observe that $h(z, 0) = f(z)$. Moreover $|\nabla_M f(z)|^2 \leq |\nabla_N h(z, 0)|^2$.

The function h is harmonic in N because

$$\Delta_N h(z, t) = \sum_{i=1}^{k_L} \left[\beta_i e^{\sqrt{\lambda_i} t} \Delta_M \phi_i(z) + \beta_i \phi_i(z) \Delta_{\mathbb{R}}(e^{\sqrt{\lambda_i} t}) \right] = 0.$$

For the harmonic extension, we don't have the mean-value property, because it is not true for all manifolds (only for the harmonic manifolds, see [17] for a complete characterization of them). What is always true is a "submean value property" (with a uniform constant) for positive subharmonic functions, see for example [15, Chapter II, Section 6]).

Observe that since h is harmonic on N , $|h|^2$ is a positive subharmonic function on N . In fact, $|h|^p$ is subharmonic for all $p \geq 1$ (for a proof see [3, Proposition 1]). Therefore, we know that for all $r < R_0(M)$

$$|h(z, t)|^2 \lesssim \int_{B(z, r/\sqrt{L}) \times I_r(t)} |h(w, s)|^2 dV(w) ds,$$

where $R_0(M) > 0$ is the injectivity radius of M and $I_r(t) = (t - r/\sqrt{L}, t + r/\sqrt{L})$. In particular,

$$(3) \quad |f(z)|^2 \leq C_r L^{(m+1)/2} \int_{B(z, r/\sqrt{L}) \times I_r} |h(w, s)|^2 dV(w) ds,$$

where $I_r = I_r(0)$.

The following result relates the L^2 -norm of f and h .

Proposition 1. *Let $r > 0$ fixed. If $f \in E_L$ then*

$$(4) \quad 2re^{-2r} \|f\|_2^2 \leq \sqrt{L} \|h\|_{L^2(M \times I_r)}^2 \leq 2re^{2r} \|f\|_2^2.$$

Therefore, for $r < R_0(M)$

$$\frac{\sqrt{L}}{2r} \|h\|_{L^2(M \times I_r)}^2 \approx \|f\|_2^2$$

with constants depending only on the manifold M .

Proof. Using the orthogonality of $\{\phi_i\}_i$ we have

$$\begin{aligned} \|h\|_{L^2(M \times I_r)}^2 &= \int_{I_r} \int_M \left| \sum_{i=1}^{k_L} \beta_i \phi_i(z) e^{\sqrt{\lambda_i} t} \right|^2 dV(z) dt \\ &= \int_{I_r} \sum_{i=1}^{k_L} \int_M |\beta_i|^2 |\phi_i(z)|^2 dV(z) e^{2\sqrt{\lambda_i} t} dt \leq \int_{I_r} e^{2\sqrt{L} t} dt \|f\|_2^2. \end{aligned}$$

Similarly, one can prove the left hand side inequality of (4). $\square$

We recall now a result proved by Schoen and Yau that estimates the gradient of harmonic functions.

Theorem (Schoen-Yau). *Let N be a complete Riemannian manifold with Ricci curvature bounded below by $-(n-1)K$ (n is the dimension of N and K a positive constant). Suppose B_a is a geodesic ball in N with radius a and h is an harmonic function on B_a . Then*

$$(5) \quad \sup_{B_{a/2}} |\nabla h| \leq C_n \left(\frac{1 + a\sqrt{K}}{a} \right) \sup_{B_a} |h|,$$

where C_n is a constant depending only on the dimension of N .

For a proof see [15, Corollary 3.2., page 21].

Remark 1. We will use Schoen and Yau's estimate in the following context. Take $N = M \times \mathbb{R}$. Observe that $\text{Ricc}(N) = \text{Ricc}(M)$ which is bounded below because M is compact. Note that N is complete because it is a product of two complete manifolds. We will take $a = r/\sqrt{L}$ ($r < R_0(M)$) and $B_a = B(z, r/\sqrt{L}) \times I_r$ (note that this is not the ball of center $(z, 0) \in N$ and radius $r/\sqrt{L}$, but it contains and it is contained in such ball of comparable radius).

Using Schoen and Yau's theorem, we deduce the global Bernstein inequality for a single eigenfunction.

Corollary 1 (Bernstein inequality). *If u is an eigenfunction of eigenvalue λ , then*

$$(6) \quad \|\nabla u\|_\infty \lesssim \sqrt{\lambda} \|u\|_\infty.$$

Proof. The harmonic extension of u is $h(z, t) = u(z)e^{\sqrt{\lambda}t}$. Applying inequality (5) to h (taking $a = R_0(M)/(2\sqrt{\lambda})$),

$$|\nabla u(z)| \lesssim \sqrt{\lambda} \|h\|_{L^\infty(M \times I_{R_0/2})} \approx \sqrt{\lambda} \|u\|_\infty.$$

□

We conjecture that in inequality (6), one can replace u by the functions $f \in E_L$, i.e.

$$\|\nabla f\|_\infty \lesssim \sqrt{L} \|f\|_\infty.$$

For instance, as a direct consequence of Green's formula, we have the L^2 -Bernstein inequality for the space E_L :

$$\|\nabla f\|_2 \lesssim \sqrt{L} \|f\|_2 \quad \forall f \in E_L.$$

For our purpose, it is sufficient a weaker Bernstein type inequality that compares the L^∞ norm of the gradient with the L^2 norm of the function.

Proposition 2. *Let $f \in E_L$. Then there exists a universal constant C such that*

$$\|\nabla f\|_\infty \leq C\sqrt{k_L}\sqrt{L} \|f\|_2.$$

For the proof, we need the following lemma.

Lemma 1. For all $f \in E_L$ and $0 < r < R_0(M)/2$,

$$|\nabla f(z)|^2 \leq C_r L^{(m+2+1)/2} \int_{B(z, r/\sqrt{L}) \times I_r} |h(w, s)|^2 dV(w) ds.$$

Proof. Using inequality (5) and the submean-value inequality for $|h|^2$, we have

$$\begin{aligned} |\nabla f(z)|^2 &\leq |\nabla h(z, 0)|^2 \lesssim \frac{L}{r^2} \sup_{B(z, r/\sqrt{L}) \times I_r} |h(w, t)|^2 \\ &\lesssim \frac{L^{(m+1+2)/2}}{\tilde{r}^{m+2+1}} \int_{B(z, \tilde{r}/\sqrt{L}) \times I_{\tilde{r}}} |h(\xi, s)|^2 dV(\xi) ds, \end{aligned}$$

where $\tilde{r} = 2r$. □

Proposition 2. Using Lemma 1, given $0 < r < R_0(M)/2$ there exists a constant C_r such that

$$|\nabla f(z)|^2 \leq C_r k_L L \sqrt{L} \int_{M \times I_r} |h(w, s)|^2 dV(w) ds \stackrel{\text{Proposition 1}}{\approx} C_r k_L L \|f\|_2^2.$$

Taking $r = R_0(M)/4$, we get that $|\nabla f(z)|^2 \leq C k_L L \|f\|_2^2$ for all $z \in M$. □

3. CHARACTERIZATION OF CARLESON MEASURES

Definition 1. Let $\mu = \{\mu_L\}_{L \geq 0}$ be a sequence of measures on M . We say that μ is an L^2 -**Carleson sequence** for M if there exists a positive constant C such that for all L and $f_L \in E_L$

$$\int_M |f_L|^2 d\mu_L \leq C \int_M |f_L|^2 dV.$$

Theorem 3. Let μ be a sequence of measures on M . Then, μ is L^2 -Carleson for M if and only if there exists a $C > 0$ such that for all L

$$(7) \quad \sup_{\xi \in M} \mu_L(B(\xi, 1/\sqrt{L})) \leq \frac{C}{k_L}.$$

Remark 2. Condition (7) can be viewed as

$$\sup_{\xi \in M} \frac{\mu_L(B(\xi, 1/\sqrt{L}))}{\text{vol}(B(\xi, 1/\sqrt{L}))} \lesssim 1.$$

First, we prove the following result.

Lemma 2. Let μ be a sequence of measures on M . Then, the following conditions are equivalent.

(1) There exists a constant $C = C(M) > 0$ such that for each L

$$\sup_{\xi \in M} \mu_L(B(\xi, 1/\sqrt{L})) \leq \frac{C}{k_L}.$$

- (2) *There exist $c = c(M) > 0$ ($c < 1$ small) and $C = C(M) > 0$ such that for all L*

$$\sup_{\xi \in M} \mu_L(B(\xi, c/\sqrt{L})) \leq \frac{C}{k_L}.$$

Proof. Obviously, the first condition implies the second one since

$$B(\xi, c/\sqrt{L}) \subset B(\xi, 1/\sqrt{L}).$$

Let's see the converse. The manifold M is covered by the union of balls of center $\xi \in M$ and radius $c/\sqrt{L}$. Taking into account the 5- r covering lemma (see [13, Chapter 2, page 23] for more details), we get a family of disjoint balls, denoted by $B_i = B(\xi_i, c/\sqrt{L})$, such that M is covered by the union of $5B_i$. This union may be finite or countable. Let $\xi \in M$ and consider $B := B(\xi, 1/\sqrt{L})$. Let n be the number of balls $\bar{B}_i$ such that $\bar{B} \cap 5\bar{B}_i \neq \emptyset$. Since $\bar{B}$ is compact, we have a finite number of these balls, so that

$$\bar{B} \subset \cup_{i=1}^n 5\bar{B}_i.$$

We claim that n is independent of L . Hence we will get our statement because

$$\mu_L(B) \leq \sum_{i=1}^n \mu_L(B(\xi_i, 5c/\sqrt{L})) \lesssim \frac{n}{k_L}.$$

We only need to check that n is independent of L . Using the triangle inequality, for all $i = 1, \dots, n$

$$B(\xi_i, c/\sqrt{L}) \subset B(\xi, 10/\sqrt{L}).$$

Therefore,

$$\cup_{i=1}^n B(\xi_i, c/\sqrt{L}) \subset B(\xi, 10/\sqrt{L}),$$

where the union is a disjoint union of balls. Now,

$$\frac{10^m}{L^{m/2}} \approx \text{vol}(B(\xi, 10/\sqrt{L})) \geq \sum_{i=1}^n \text{vol}(B_i) \approx n \frac{c^m}{L^{m/2}}.$$

Hence, $n \lesssim (10/c)^m$ and we can choose it independently of L . $\square$

Now we can prove the characterization of the Carleson measures.

Theorem 3. Assume condition (7). We need to prove the existence of a constant $C > 0$ (independent of L) such that for each $f \in E_L$

$$\int_M |f|^2 d\mu_L \leq C \int_M |f|^2 dV.$$

Let $f \in E_L$ with L and $r > 0$ (small) fixed.

$$\begin{aligned}
\int_M |f(z)|^2 d\mu_L &\stackrel{(3)}{\leq} C_r L^{(m+1)/2} \int_M \int_{B(z, r/\sqrt{L}) \times I_r} |h(w, s)|^2 dV(w) ds d\mu_L(z) \\
&= C_r L^{(m+1)/2} \int_{M \times I_r} |h(w, s)|^2 \mu_L(B(w, r/\sqrt{L})) dV(w) ds \\
&\leq C_r L^{(m+1)/2} \frac{1}{k_L} \int_{M \times I_r} |h(w, s)|^2 dV(w) ds \stackrel{\text{Proposition 1}}{\approx} \|f\|_2^2
\end{aligned}$$

with constants independent of L . Therefore, $\mu = \{\mu_L\}_L$ is L^2 -Carleson for M .

For the converse, assume that μ is L^2 -Carleson for M . We have to show the existence of a constant C such that for all $L \geq 1$ and $\xi \in M$, $\mu_L(B(\xi, c/\sqrt{L})) \leq C/k_L$ (for some small constant $c > 0$). We will argue by contradiction, i.e. assume that for all $n \in \mathbb{N}$ there exists L_n and a ball B_n of radius $c/\sqrt{L_n}$ such that $\mu_{L_n}(B_n) > n/k_{L_n} \approx n/L_n^{m/2}$ (c will be chosen later). Let b_n be the center of the ball B_n . Define $F_n(w) = K_{L_n}(b_n, w)$. Note that the function $L_n^{-m/4} F_n \in E_{L_n}$ and $\|F_n\|_2^2 = K_{L_n}(b_n, b_n) \approx L_n^{m/2}$. Therefore,

$$\begin{aligned}
C &\approx \int_M |L_n^{-m/4} F_n|^2 dV \gtrsim \int_M |L_n^{-m/4} F_n|^2 d\mu_{L_n} \gtrsim \int_{B_n} |L_n^{-m/4} F_n|^2 d\mu_{L_n} \\
&\geq \inf_{w \in B_n} |L_n^{-m/4} F_n(w)|^2 \mu_{L_n}(B_n) \gtrsim \inf_{w \in B_n} |F_n(w)|^2 \frac{n}{L_n^m}.
\end{aligned}$$

Now we will study this infimum. Let $w \in B_n = B(b_n, c/\sqrt{L_n})$. Then

$$|F_n(b_n)| - |F_n(w)| \leq |F_n(b_n) - F_n(w)| \leq \frac{c}{\sqrt{L_n}} \|\nabla F_n\|_\infty \leq$$

$$\stackrel{\text{Proposition 2}}{\leq} \frac{c}{\sqrt{L_n}} C_1 \sqrt{k_{L_n}} \sqrt{L_n} \|F_n\|_2 \approx c C_1 k_{L_n}.$$

We pick c small enough so that

$$\inf_{B_n} |F_n(w)|^2 \geq C L_n^m.$$

Finally, we have shown that $C \gtrsim n \forall n$. This gives us the contradiction. $\square$

The following result is a Plancherel-Pólya type theorem but in the context of the Paley-Wiener spaces E_L .

Theorem 4 (Plancherel-Pólya Theorem). *Let $\mathcal{Z}$ be a triangular family of points in M , i.e. $\mathcal{Z} = \{z_{Lj}\}_{j \in \{1, \dots, m_L\}, L \geq 1} \subset M$. Then $\mathcal{Z}$ is a finite union of uniformly separated families, if and only if there exists*

a constant $C > 0$ such that for all $L \geq 1$ and $f_L \in E_L$

$$(8) \quad \frac{1}{k_L} \sum_{j=1}^{m_L} |f_L(z_{Lj})|^2 \leq C \int_M |f_L(\xi)|^2 dV(\xi).$$

Remark 3. The above result is interesting because the inequality (8) means that the sequence of normalized reproducing kernels is a Bessel sequence for E_L , i.e.

$$\sum_{j=1}^{m_L} |\langle f, \tilde{K}_L(\cdot, z_{Lj}) \rangle|^2 \lesssim \|f\|_2^2 \quad \forall f \in E_L,$$

where $\{\tilde{K}_L(\cdot, z_{Lj})\}_j$ are the normalized reproducing kernels. Note that $|\tilde{K}_L(\cdot, z_{Lj})|^2 \approx |K_L(\cdot, z_{Lj})|^2 k_L^{-1}$. That's the reason why the quantity k_L appears in the inequality (8).

Proof. This is a consequence of Theorem 3 applied to the measures

$$\mu_L = \frac{1}{k_L} \sum_{j=1}^{m_L} \delta_{z_{Lj}}, \quad L \geq 1.$$

□

4. CHARACTERIZATION OF LOGVINENKO-SEREDA SETS

Before we state the characterization, we would like to recall some history of these kind of inequalities. The classical Logvinenko-Sereda (L-S) theorem describes some equivalent norms for functions in the Paley-Wiener space PW_Ω^p . The precise statements is the following:

Theorem 5 (L-S). *Let Ω be a bounded set and $1 \leq p < +\infty$. A set $E \subset \mathbb{R}^d$ satisfies*

$$\int_{\mathbb{R}^d} |f(x)|^p dx \leq C_p \int_E |f(x)|^p dx, \quad \forall f \in PW_\Omega^p,$$

if and only if there is a cube $K \subset \mathbb{R}^d$ such that

$$\inf_{x \in \mathbb{R}^d} |(K+x) \cap E| > 0.$$

One can find the original proof in [9] and another proof can be found in [4, p. 112-116].

Luecking in [10] studied this notion for the Bergman spaces. Following his ideas, in [12], it has been proved the following result.

Theorem 6. *Let $1 \leq p < +\infty$, μ be a doubling measure and let $\mathcal{E} = \{E_L\}_{L \geq 0}$ be a sequence of sets in $\mathbb{S}^d$. Then $\mathcal{E}$ is $L^p(\mu)$ -L-S if and only if $\mathcal{E}$ is μ -relatively dense.*

For the precise definitions and notations see [12]. Using the same ideas, we will prove the above theorem for the case of our arbitrary compact manifold M and the measure given by the volume element.

In what follows, $\mathcal{A} = \{A_L\}_L$ will be a sequence of sets in M .

Definition 2. We say that $\mathcal{A}$ is L-S if there exists a constant $C > 0$ such that for any L and $f_L \in E_L$

$$\int_M |f_L|^2 dV \leq C \int_{A_L} |f_L|^2 dV.$$

Definition 3. The sequence of sets $\mathcal{A} \subset M$ is relatively dense if there exists $r > 0$ and $\rho > 0$ such that for all L

$$\inf_{z \in M} \frac{\text{vol}(A_L \cap B(z, r/\sqrt{L}))}{\text{vol}(B(z, r/\sqrt{L}))} \geq \rho > 0.$$

Remark 4. It is equivalent to have this property for all $L \geq L_0$ for some L_0 fixed.

Our main statement is the following:

Theorem 7. $\mathcal{A}$ is L-S if and only if $\mathcal{A}$ is relatively dense.

We shall prove the two implications in the statement separately. First we will show that this condition is necessary. Before proceeding, we construct functions in E_L with a desired decay of its L^2 -integral outside a ball.

Proposition 3. Given $\xi \in M$ and $\epsilon > 0$, there exist functions $f_L = f_{L,\xi} \in E_L$ and $R_0 = R_0(\epsilon, M) > 0$ such that

- (1) $\|f_L\|_2 = 1$.
- (2)

$$\int_{M \setminus B(\xi, R_0/\sqrt{L})} |f_L|^2 dV < \epsilon \quad \forall L.$$

- (3) For all $L \geq 1$ and any subset $A \subset M$,

$$\int_A |f_L|^2 dV \leq C_1 \frac{\text{vol}(A \cap B(\xi, R_0/\sqrt{L}))}{\text{vol}(B(\xi, R_0/\sqrt{L}))} + \epsilon,$$

where C_1 is a constant independent of L , ξ and f_L .

Remark. In the above Proposition, the R_0 does not depend on the point ξ .

Proof. Given $z, \xi \in M$ and $L \geq 1$, let $S_L^N(z, \xi)$ denote the Riesz kernel of index $N \in \mathbb{N}$ associated to the Laplacian, i.e

$$S_L^N(z, \xi) = \sum_{i=1}^{k_L} \left(1 - \frac{\lambda_i}{L}\right)^N \phi_i(z) \phi_i(\xi).$$

Note that $S_L^0(z, \xi) = K_L(z, \xi)$. The Riesz kernel satisfies the following inequality.

$$(9) \quad |S_L^N(z, \xi)| \leq CL^{m/2}(1 + \sqrt{L}d(z, \xi))^{-N-1}.$$

This estimate has its origins in Hörmander's article [6, Theorem 5.3]. Estimate (9) can be found also in [16, Lemma 2.1].

Note that on the diagonal, $S_L^N(z, z) \approx C_N L^{m/2}$. The upper bound is trivial by the definition and the lower bound follows from

$$S_L^N(z, z) \geq \sum_{i=1}^{k_{L/2}} \left(1 - \frac{\lambda_i}{L}\right)^N \phi_i(z) \phi_i(z) \geq 2^{-N} K_{L/2}(z, z) \approx C_N L^{m/2}.$$

Similarly we observe that $\|S_L^N(\cdot, \xi)\|_2^2 \approx C_N L^{m/2}$.

Given $\xi \in M$, define for all $L \geq 1$

$$f_{L,\xi}(z) := f_L(z) = \frac{S_L^N(z, \xi)}{\|S_L^N(\cdot, \xi)\|_2}.$$

We will choose the order N later. Each f_L belongs to the space E_L and has unit L^2 -norm. Let us verify the second property claimed in Proposition 3. Fix a radius R . Using the estimate (9),

$$\int_{M \setminus B(\xi, R/\sqrt{L})} |f_L|^2 dV \leq C_N L^{m/2} \int_{M \setminus B(\xi, R/\sqrt{L})} \frac{dV}{(\sqrt{L}d(z, \xi))^{2(N+1)}} = (\star)$$

For any $t \geq 0$, consider the following set.

$$A_t := \left\{ z \in M : d(z, \xi) \geq \frac{R}{\sqrt{L}}, \quad d(z, \xi) < \frac{t^{-1/(2(N+1))}}{\sqrt{L}} \right\}.$$

Note that for $t > R^{-2(N+1)}$, $A_t = \emptyset$ and for $t < R^{-2(N+1)}$, $A_t \subset B(\xi, t^{-1/(2(N+1))}/\sqrt{L})$. Using the distribution function, we have:

$$(\star) = C_N L^{m/2} \int_0^{R^{-2(N+1)}} \text{vol}(A_t) dt \leq C_N \frac{1}{R^{2(N+1)-m}},$$

provided $N + 1 > m/2$. Thus if we pick R_0 big enough we get

$$(10) \quad \int_{M \setminus B(\xi, R_0/\sqrt{L})} |f_L|^2 dV < \epsilon.$$

Now the third property claimed in Proposition 3 follows from (10). Indeed, given any subset A in the manifold M ,

$$\int_A |f_L|^2 dV \leq \int_{A \cap B(\xi, R_0/\sqrt{L})} |f_L|^2 dV + \epsilon.$$

Observe that

$$\begin{aligned} \int_{A \cap B(\xi, R_0/\sqrt{L})} |f_L|^2 dV &\lesssim C_N L^{m/2} \int_{A \cap B(\xi, R_0/\sqrt{L})} \frac{dV(z)}{(1 + \sqrt{L}d(z, \xi))^{2(N+1)}} \\ &\lesssim C_N R_0^m \frac{\text{vol}(A \cap B(\xi, R_0/\sqrt{L}))}{\text{vol}(B(\xi, R_0/\sqrt{L}))}. \end{aligned}$$

□

Now we are ready to prove one of the implications in the characterization of the L-S sets.

Proposition 4. *Assume $\mathcal{A}$ is L-S. Then it is relatively dense.*

Proof. Assume $\mathcal{A}$ is L-S, i.e.

$$\int_M |f_L|^2 dV \leq C \int_{A_L} |f_L|^2 dV.$$

Let $\xi \in M$ be an arbitrary point. Fix $\epsilon > 0$ and consider the R_0 and the functions $f_L \in E_L$ given by Proposition 3. Using the third property of Proposition 3 for the sets A_L , we get for all $L \geq 1$,

$$1 = \|f_L\|_2^2 \leq C \int_{A_L} |f_L|^2 \leq CC_1 \frac{\text{vol}(A_L \cap B(\xi, R_0/\sqrt{L}))}{\text{vol}(B(\xi, R_0/\sqrt{L}))} + C\epsilon,$$

where C_1 is a constant independent of L , ξ and f_L . Therefore, we have proved that there exist constants c_1 and c_2 such that

$$\frac{\text{vol}(A_L \cap B(\xi, R_0/\sqrt{L}))}{\text{vol}(B(\xi, R_0/\sqrt{L}))} \geq c_1 - c_2\epsilon.$$

Hence, $\mathcal{A}$ is relatively dense provided $\epsilon > 0$ is small enough. □

Before we continue, we will prove a result concerning the uniform limit of harmonic functions with respect to some metric.

Lemma 3. *Let $\{H_n\}_n$ be a family of uniformly bounded real functions defined on the ball $\mathbb{B}(0, \rho) \subset \mathbb{R}^d$ for some $\rho > 0$. Let g be a non-singular $\mathcal{C}^\infty$ metric such that g and all its derivatives are uniformly bounded and $g_{ij}(0) = \delta_{ij}$. Define $g_n(z) = g(z/L_n)$ (the rescaled metrics) where L_n is a sequence of values tending to ∞ as n increases. Assume that the family $\{H_n\}_n$ converges uniformly on compact subsets of $\mathbb{B}(0, \rho)$ to a limit function $H : \mathbb{B}(0, \rho) \rightarrow \mathbb{R}$ and H_n is harmonic with respect to the metric g_n (i.e. $\Delta_{g_n} H_n = 0$). Then, the limit function H is harmonic in the Euclidean sense.*

Proof. Let $\varphi \in \mathcal{C}_c^\infty(\mathbb{B}(0, \rho))$. We have

$$\int_{\mathbb{B}(0, \rho)} \Delta_g f \varphi dV = \int_{\mathbb{B}(0, \rho)} f \Delta_g \varphi dV.$$

It is a direct computation to see that $\Delta_{g_n}\varphi \rightarrow \Delta\varphi$ uniformly and $\Delta_{g_n}\varphi$ is uniformly bounded on $\mathbb{B}(0, \rho)$. Then

$$0 = \int_{\mathbb{B}(0, \rho)} H_n \Delta_{g_n} \varphi dV_{g_n} \rightarrow \int_{\mathbb{B}(0, \rho)} H \Delta \varphi dm(z) = \int_{\mathbb{B}(0, \rho)} \Delta H \varphi dm(z).$$

Therefore, the limit function H is harmonic in the weak sense. Applying Weyl's lemma, H is harmonic in the Euclidean sense. $\square$

Remark 5. The above argument also holds if we have a sequence of metrics g_n converging uniformly to g whose derivatives also converge uniformly to the derivatives of g . In this case, the conclusion would be that the limit is harmonic with respect to the limit metric g .

Now, we shall prove the sufficient condition of the main result.

Proposition 5. *If $\mathcal{A}$ is relatively dense then it is L -S.*

Proof. Fix $\epsilon > 0$ and $r > 0$. Let $D := D_{\epsilon, r, f_L}$ be

$$D = \{z \in M : |f_L(z)|^2 = |h_L(z, 0)|^2 \geq \epsilon \int_{B(z, \frac{r}{\sqrt{L}}) \times I_r} |h_L(\xi, t)|^2 dV(\xi) dt\}.$$

The norm of f_L is almost concentrated on D because

$$\begin{aligned} \int_{M \setminus D} |f_L(z)|^2 dV(z) &\lesssim \\ &\lesssim \epsilon \frac{1}{l(I_r)} \int_{M \times I_r} |h_L(\xi, t)|^2 \frac{L^{m/2}}{r^m} \int_{(M \setminus D) \cap B(\xi, r/\sqrt{L})} dV(z) dV(\xi) dt \\ &\lesssim \epsilon \frac{1}{l(I_r)} \int_{M \times I_r} |h_L(\xi, t)|^2 dV(\xi) dt \stackrel{\text{Proposition 1}}{\lesssim} e^{2r} \epsilon \int_M |f_L|^2 dV. \end{aligned}$$

It is enough to prove

$$(11) \quad \int_D |f_L|^2 dV \lesssim \int_{A_L} |f_L|^2$$

with constants independent of L and for this, it is sufficient to show that there exists a constant $C > 0$ such that for all $w \in D$

$$(12) \quad |f_L(w)|^2 \leq \frac{C}{\text{vol}(B(w, r/\sqrt{L}))} \int_{A_L \cap B(w, r/\sqrt{L})} |f_L(\xi)|^2 dV(\xi).$$

Because then, (11) follows by integrating (12) over D . So we need to prove (12). Assume it is not true. This means that for all $n \in \mathbb{N}$ there exists L_n , functions $f_n \in E_{L_n}$ and $w_n \in D_n = D_{\epsilon, r, f_n}$ such that

$$|f_n(w_n)|^2 > \frac{n}{\text{vol}(B(w_n, r/\sqrt{L_n}))} \int_{A_{L_n} \cap B(w_n, r/\sqrt{L_n})} |f_n|^2 dV.$$

By the compactness of M , there exists $\rho_0 = \rho_0(M) > 0$ such that for all $w \in M$, the exponential map, $\exp_w : \mathbb{B}(0, \rho_0) \rightarrow B(w, \rho_0)$, is

a diffeomorphism and $(B(w, \rho_0), \exp_w^{-1})$ is a normal coordinate chart, where w is mapped to 0 and the metric g verifies $g_{ij}(0) = \delta_{ij}$.

For all $n \in \mathbb{N}$, take $\exp_n(z) := \exp_{w_n}(rz/\sqrt{L_n})$ which is defined in $\mathbb{B}(0, 1)$ and act as follows:

$$\begin{aligned} \exp_n : \mathbb{B}(0, 1) &\longrightarrow \mathbb{B}(0, r/\sqrt{L_n}) \longrightarrow B(w_n, r/\sqrt{L_n}) \\ z &\longrightarrow \frac{rz}{\sqrt{L_n}} \longrightarrow \exp_{w_n}(rz/\sqrt{L_n}) =: w \end{aligned}$$

Consider $F_n(z) := f_n(\exp_n(z)) : \mathbb{B}(0, 1) \rightarrow B(w_n, r/\sqrt{L_n}) \xrightarrow{f_n} \mathbb{R}$ and the corresponding harmonic extension h_n of f_n . Set

$$H_n(z, t) := h_n(\exp_n(z), rt/\sqrt{L_n}),$$

defined on $\mathbb{B}(0, 1) \times J_1$ (where $J_1 = (-1, 1)$).

Let μ_n be the measure such that $d\mu_n(z) = \sqrt{|g|(\exp_{w_n}(rz/\sqrt{L_n}))} dm(z)$. Note that

$$\int_{B(w_n, r/\sqrt{L_n})} |f_n|^2 dV = \frac{r^m}{L_n^{m/2}} \int_{\mathbb{B}(0, 1)} |F_n(z)|^2 d\mu_n(z).$$

Therefore, we have

$$\int_{B(w_n, r/\sqrt{L_n})} |f_n|^2 dV \approx \int_{\mathbb{B}(0, 1)} |F_n|^2 d\mu_n.$$

We will normalize H_n so that

$$\int_{\mathbb{B}(0, 1) \times J_1} |H_n(w, s)|^2 d\mu_n(w) ds = 1.$$

As $w_n \in D_n$, we have

$$\begin{aligned} |F_n(0)|^2 &= |f_n(w_n)|^2 \geq \epsilon \int_{B(w_n, r/\sqrt{L_n}) \times I_r} |h_n(w, t)|^2 dV dt \\ &\approx \epsilon \int_{\mathbb{B}(0, 1) \times J_1} |H_n(w, s)|^2 d\mu_n(w) ds = \epsilon. \end{aligned}$$

Since $|h_n|^2$ is subharmonic,

$$|F_n(0)|^2 = |h_n(w_n, 0)|^2 \lesssim \int_{\mathbb{B}(0, 1) \times J_1} |H_n(w, s)|^2 d\mu_n(w) ds = 1.$$

Hence, we have $\forall n \in \mathbb{N} \quad 0 < \epsilon \lesssim |F_n(0)|^2 \lesssim 1$.

Using the assumption,

$$\frac{1}{n} \gtrsim \frac{1}{\text{vol}(B(w_n, r/\sqrt{L_n}))} \int_{A_{L_n} \cap B(w_n, r/\sqrt{L_n})} |f_n|^2 dV \approx \int_{B_n \cap \mathbb{B}(0, 1)} |F_n|^2 d\mu_n,$$

where B_n is such that $\exp_n(B_n \cap \mathbb{B}(0, 1)) = A_{L_n} \cap B(w_n, r/\sqrt{L_n})$. So we have that

$$\begin{cases} \forall n \ 0 < \epsilon \lesssim |F_n(0)|^2 \lesssim 1 \\ \forall n \ \int_{\mathbb{B}(0,1) \cap B_n} |F_n|^2 d\mu_n \lesssim \frac{1}{n} \end{cases}.$$

In fact, $0 < \epsilon \lesssim |H_n(0, 0)|^2 \lesssim 1$ (by the definition) and one can prove that $|H_n|^2 \lesssim 1$. Indeed, if $(z, s) \in \mathbb{B}(0, 1/2) \times J_{1/2}$, let $w = \exp_n(z) \in B(w_n, r/(2\sqrt{L_n}))$ and $t = rs/\sqrt{L_n} \in I_{r/2}$. Then

$$\begin{aligned} |H_n(z, s)|^2 &= |h_n(w, t)|^2 \lesssim \int_{B(w, r/(2\sqrt{L_n})) \times I_{r/2}(t)} |h_n|^2 \\ &\lesssim \int_{B(w_n, r/\sqrt{L_n}) \times I_r} |h_n|^2 dV dt = 1. \end{aligned}$$

Therefore, working with $1/2$ instead of 1 we have $|H_n|^2 \lesssim 1$ for all n .

The sequence $\{H_n\}_n$ is equicontinuous in $\mathbb{B}(0, 1) \times J_1$. Indeed, consider $(w, t) \in B(w_n, r/(4\sqrt{L_n})) \times I_{r/4}$ and $(\tilde{w}, \tilde{t}) \in B(w, \tilde{r}r/\sqrt{L_n}) \times I_{\tilde{r}r}(t)$, then there exists some small $\delta > 0$ such that

$$|h_n(w, t) - h_n(\tilde{w}, \tilde{t})| \leq \frac{\tilde{r}}{\sqrt{L_n}} r \sup_{B(w, \delta/\sqrt{L_n}) \times I_\delta(t)} |\nabla h_n| \leq (\star)$$

Taking $\tilde{r}$ small enough so that $\delta \leq r/4$ and using Schoen and Yau's estimate (5), we have

$$(\star) \leq \frac{\tilde{r}r}{\sqrt{L_n}} \sup_{B(w_n, r/(2\sqrt{L_n})) \times I_{r/2}} |\nabla h_n| \lesssim \frac{\tilde{r}r}{\sqrt{L_n}} \frac{1}{\frac{r}{\sqrt{L_n}}} \sup_{B(w_n, r/\sqrt{L_n}) \times I_r} |h_n| \lesssim \tilde{r}.$$

So we have proved that $|h_n(w, t) - h_n(\tilde{w}, \tilde{t})| \leq C\tilde{r}$. Take $\tilde{r}$ small enough so that $C\tilde{r} < \epsilon$. Let $(z, s) \in \mathbb{B}(0, 1/4) \times J_{1/4}$ and $(\tilde{z}, \tilde{s}) \in \mathbb{B}(z, \tilde{r}) \times (s - \tilde{r}, s + \tilde{r})$. Consider $w = \exp_n(z)$, $t = rs/\sqrt{L_n}$, $\tilde{w} = \exp_n(\tilde{z})$ and $\tilde{t} = r\tilde{s}/\sqrt{L_n}$. We have proved that for all $\epsilon > 0$ there exists $\tilde{r} > 0$ (small) such that for all $(z, s) \in \mathbb{B}(0, 1/4) \times J_{1/4}$:

$$|H_n(z, s) - H_n(\tilde{z}, \tilde{s})| < \epsilon \quad \text{if } |z - \tilde{z}| < \tilde{r}, \ |s - \tilde{s}| < \tilde{r} \quad \forall n.$$

Change $1/4$ by 1 . So the sequence H_n is equicontinuous.

Hence the family $\{H_n\}_n$ is equicontinuous and uniformly bounded on $\mathbb{B}(0, 1) \times J_1$. Therefore, by Ascoli-Arzelà's theorem there exists a partial sequence (denoted as the sequence itself) such that $H_n \rightarrow H$ uniformly on compact subsets of $\mathbb{B}(0, 1) \times J_1$. Since $F_n(z) = H_n(z, 0)$, we get a function $F(z) := H(z, 0) : \mathbb{B}(0, 1) \rightarrow \mathbb{R}$ which is the limit of F_n (uniformly on compact subsets of $\mathbb{B}(0, 1)$).

By hypothesis, the sequence $\{A_L\}_L$ is relatively dense. Taking into

account that $\text{vol}(B(w_n, r/\sqrt{L_n})) = \frac{r^m}{L_n^{m/2}} \mu_n(\mathbb{B}(0, 1))$, we get that

$$(13) \quad \inf_n \mu_n(B_n) \geq \rho > 0,$$

where we have denoted $B_n \cap \mathbb{B}(0, 1)$ by B_n .

Let τ_n be such that $d\tau_n = \chi_{B_n} d\mu_n$. From a standard argument (τ_n are supported in a ball), we know the existence of a weak $*$ -limit of a subsequence of τ_n , denoted by τ . This subsequence will be noted as the sequence itself. Using (13), we know that τ is not identically 0. Now we have that

$$\int_{\mathbb{B}(0,1)} |F|^2 d\tau = 0.$$

Therefore, $F = 0$ τ -a.e. in $\mathbb{B}(0, 1)$. Now for all $K \subset \mathbb{B}(0, 1)$ compact

$$\int_K |F|^2 d\tau = 0,$$

therefore $F = 0$ in $\text{supp}\tau$. Let $\overline{\mathbb{B}(a, s)} \subset \mathbb{B}(0, 1)$ such that $\overline{\mathbb{B}(a, s)} \cap \text{supp}\tau \neq \emptyset$, then using the fact $B_n \subset \mathbb{B}(0, 1)$,

$$\tau_n(\overline{\mathbb{B}(a, s)}) \leq \int_{\overline{\mathbb{B}(a, s)}} d\mu_n \approx \frac{L_n^{m/2}}{r^m} \text{vol}(B(\exp_n(a), sr/\sqrt{L_n})) \approx s^m.$$

Therefore $\tau_n(\overline{\mathbb{B}(a, s)}) \lesssim s^m$ for all n . Hence, in the limit $\tau(\overline{\mathbb{B}(a, s)}) \lesssim s^m$. In short,

(1) We have sets $B_n \subset \mathbb{B}(0, 1)$ such that

$$\rho \leq \mu_n(B_n) \leq \mu_n(\mathbb{B}(0, 1)) \approx 1.$$

(2) We have measures τ_n weakly- $*$ converging to τ (not identically 0).

(3) $\tau(\overline{\mathbb{B}(a, s)}) \lesssim s^m$ for all $\overline{\mathbb{B}(a, s)} \subset \mathbb{B}(0, 1)$.

(4) $|F| = 0$ τ -a.e. in $\mathbb{B}(0, 1)$.

(5) $|F(0)| > 0$ and $|F| \lesssim 1$.

Assume we know that H is real analytic, then $F(z)$ is real analytic. Federer ([2, Theorem 3.4.8]) proved that the $(m-1)$ -Hausdorff measure $\mathcal{H}^{m-1}(F^{-1}(0)) < \infty$. Hence $\mathcal{H}^{m-1}(\text{supp}\tau) \leq \mathcal{H}^{m-1}(F^{-1}(0)) < \infty$. This implies that the Hausdorff dimension $\dim_{\mathcal{H}}(\text{supp}\tau) \leq m-1$. On the other hand, since $\tau(\overline{\mathbb{B}(a, s)}) \lesssim s^m$, we have

$$0 < \tau(\text{supp}\tau) \lesssim \mathcal{H}^m(\text{supp}\tau)$$

and this implies that $\dim_{\mathcal{H}}(\text{supp}\tau) \geq m$ by Frostman's lemma. So we reach to a contradiction and the proof is complete. We only need to check that H is real analytic. In fact, we will show that H is harmonic. We have the following properties:

(1) Observe that the family of measure $d\mu_n$ converges uniformly to the ordinary Euclidean measure because $g_{ij}(\exp_{w_n}(rz/\sqrt{L_n})) \rightarrow$

- $g_{ij}(\exp_{w_0}(0)) = \delta_{ij}$, where w_0 is the limit point of some subsequence of w_n (recall that we are taking normal coordinate charts).
- (2) If $g_n(z) := g(rz/\sqrt{L_n})$ (i.e. g_n is the rescaled metric), then $\Delta_{(g_n, Id)} H_n(z, s) = 0$ for all $(z, s) \in \mathbb{B}(0, 1) \times J_1$, by construction.
- (3) The functions H_n are uniformly bounded and converge uniformly on compact subsets of $\mathbb{B}(0, 1) \times J_1$.

We are in the hypothesis of Lemma 3 that guarantees the harmonicity of H in the Euclidean sense. This concludes the proof of the proposition. $\square$

REFERENCES

- [1] H. Donnelly, C. Fefferman. Growth and Geometry of Eigenfunctions of the Laplacian. Analysis and partial differential equations, p. 635-655, Lecture Notes in Pure and Appl. Math., 122, Dekker, New York, 1990.
- [2] H. Federer. Geometric measure theory. Die Grundlehren der mathematischen Wissenschaften, Band 153 Springer-Verlag New York Inc., New York 1969.
- [3] R.E. Greene, H. Wu. Integrals of Subharmonic Functions on Manifolds of Non-negative Curvature. *Inventiones mathematicae*, Volume 27, 1974, p. 265-298.
- [4] V. Havin, B. Jorické. The uncertainty principle in harmonic analysis, Springer-Verlag, Berlin, 1994.
- [5] L. Hörmander. The spectral function of an elliptic operator. *Acta Math.* 121 (1968), p. 193-218.
- [6] L. Hörmander. On the Riesz means of spectral functions and eigenfunction expansions for elliptic differential operators. Some Recent Advances in the Basic Sciences, Vol. 2 (Proc. Annual Sci. Conf., Belfer Grad. School Sci., Yeshiva Univ., New York, 1965-1966), p. 155-202.
- [7] J. Jost, Riemannian geometry and geometric analysis. Fifth edition. Universitext. Springer-Verlag, Berlin, 2008. xiv+583 pp. ISBN: 978-3-540-77340-5.
- [8] Peter Li, Richard Schoen. L^p and mean value properties of subharmonic functions on Riemannian manifolds. *Acta Math.* 153 (1984), no. 3-4, p. 279-301.
- [9] V.N. Logvinenko, Ju F. Sereda. Equivalent norms in spaces of entire functions of exponential type. *Teor. funktsii, funkts. analiz i ich prilozhenia* 20, 102-111, 175, 1974.
- [10] D. H. Luecking. Equivalent norms on L^p spaces of harmonic functions, *Monatsh. Math.* 96, no. 2, 133-141, 1983.
- [11] J. Marzo. Marcinkiewicz-Zygmund inequalities and interpolation by spherical harmonics. *J. Funct. Anal.* 250 (2007), no. 2, p. 559-587.
- [12] J. Marzo, J. Ortega-Cerdà. Equivalent norms for polynomials on the sphere. *Int. Math. Res. Not. IMRN* 2008, no. 5, Art. ID rnm 154, 18p.
- [13] Pertti Mattila. Geometry of Sets and Measures in Euclidean Spaces. Fractals and Rectifiability. Cambridge University Press.
- [14] J. Ortega-Cerdà, J. Saludes. Marcinkiewicz-Zygmund inequalities. *J. Approx. Theory* 145 (2007), no. 2, p. 237-252.
- [15] R. Schoen, S.-T. Yau. Lectures on Differential Geometry. Conference Proceedings and Lecture Notes in Geometry and Topology, Volume I.
- [16] Christopher D. Sogge. On the Convergence of Riesz Means on Compact Manifolds. *The Annals of Mathematics*, Second Series, Vol. 126, no. 3, p. 439-447.
- [17] T. J. Willmore. Mean value theorems in harmonic riemannian spaces. *J. London Math. Soc.* 25, (1950), p. 54-57.

DEPT. MATEMÀTICA APLICADA I ANÀLISI, UNIVERSITAT DE BARCELONA,
GRAN VIA 585, 08007 BARCELONA, SPAIN
E-mail address: jortega@ub.edu

DEPT. MATEMÀTICA APLICADA I ANÀLISI, UNIVERSITAT DE BARCELONA,
GRAN VIA 585, 08007 BARCELONA, SPAIN
E-mail address: bharti.pridhnani@ub.edu